

Jaboticaba-Induced Endothelium-Independent Vasodilating Effect on Isolated Arteries

Daniela Medeiros Lobo de Andrade¹, Leonardo Luis Borges², Ieda Maria Sapateiro Torres¹, Edemilson Cardoso da Conceição², Matheus Lavorenti Rocha¹

Laboratório de Farmacologia Cardiovascular – Faculdade de Farmácia - Universidade Federal de Goiás (UFG)¹; Laboratório de Pesquisa em Produtos Naturais – Faculdade de Farmácia – Universidade Federal de Goiás (UFG)²; Goiânia, GO – Brazil

Abstract

Background: Despite the important biological effects of jaboticaba, its actions on the cardiovascular system have not been clarified.

Objectives: To determine the effects of jaboticaba hydroalcoholic extract (JHE) on vascular smooth muscle (VSM) of isolated arteries.

Methods: Endothelium-denuded aortic rings of rats were mounted in isolated organ bath to record isometric tension. The relaxant effect of JHE and the influence of K⁺ channels and Ca²⁺ intra- and extracellular sources on JHE-stimulated response were assessed.

Results: Arteries pre-contracted with phenylephrine showed concentration-dependent relaxation (0.380 to 1.92 mg/mL). Treatment with K⁺ channel blockers (tetraethyl-ammonium, glibenclamide, 4-aminopyridine) hindered relaxation due to JHE. In addition, phenylephrine-stimulated contraction was hindered by previous treatment with JHE. Inhibition of sarcoplasmic reticulum Ca²⁺ ATPase did not change relaxation due to JHE. In addition, JHE inhibited the contraction caused by Ca²⁺ influx stimulated by phenylephrine and KCl (75 mM).

Conclusion: JHE induces endothelium-independent vasodilation. Activation of K⁺ channels and inhibition of Ca²⁺ influx through the membrane are involved in the JHE relaxant effect. (Arq Bras Cardiol. 2016; 107(3):223-229)

Keywords: *Jaboticaba* (*Myrciaria cauliflora*); Trees; Vasodilatation; Calcium Channels; Muscle, Smooth Vascular.

Introduction

Cardiovascular diseases are a major cause of death worldwide, among which hypertension accounts for 9.4 million deaths per year.¹ Around 1 billion adults in the world have hypertension, and that figure will have increased by 25% in 10 years.²

Vascular tonus regulation is fundamental to appropriate blood pressure control. Blood vessel contraction and dilation in response to physiological demands are controlled by changes in the intracellular concentration of Ca²⁺ in vascular smooth muscle (VSM) cells. The increase in intracellular concentration of Ca²⁺ occurs via both Ca²⁺ influx through the plasma membrane and Ca²⁺ release from inner sources, such as the sarcoplasmic reticulum.^{3,4} Effective drugs to blood pressure control, such as

nifedipine, verapamil and diltiazem, which act as Ca²⁺ channel blockers, induce vasodilation and reduce blood pressure.⁵

The use of natural products as an alternative treatment for hypertension has been extensively studied, being known to induce hypotension with minimum side effects.^{6,7} *Jaboticaba* (*Myrciaria cauliflora*), also known as Brazilian grape, is a hard-skinned berry of the Myrtaceae family, largely distributed in Brazil. It can be consumed fresh or in the form of liqueurs, wines, jams and sweets, and its consumption has increased in Brazil and worldwide.^{8,9} In addition to the use of *jaboticaba* as food and beverage, in folk medicine, that fruit is used to treat some diseases, such as asthma, inflammations, and gastrointestinal and cardiovascular disorders.¹⁰ Recent findings have shown that *jaboticaba* can decrease oxidative process,¹¹ hyperglycemia associated with insulin resistance¹² and dyslipidemia.¹³ In addition, that species has a proven endothelium-dependent hypotensive and vasodilating effect, mediated by nitric oxide pathway.¹⁴

Considering that *jaboticaba* has important biological effects and that its action on the cardiovascular system has been little studied, this study was aimed at assessing the possible effect of the *jaboticaba* extract directly on the VSM, mainly its effect on Ca²⁺ influx through the plasma membrane and activation of K⁺ channels.

Mailing Address: Matheus Lavorenti Rocha •

UFG – Faculdade de Farmácia. Rua 240, esquina com 5ª Avenida, s/n, Setor Leste Universitário. Postal Code 74605-170, Goiânia, Goiás – Brazil.

E-mail: matheusroch@yahoo.com.br, rochaml@ufg.br

Manuscript received November 25, 2015; revised manuscript March 04, 2016; accepted March 07, 2016.

DOI: 10.5935/abc.20160118

Methods

Preparation of jaboticaba extract

For this study, the plant specimens were donated by the "Jaboticabal" wine house in the city of Hidrolândia, Goiás state, Brazil. A sample of the plant specimen was stored at the herbarium of the department of botany of the Federal University of Goiás (UFG), Goiânia, Goiás state, Brazil (n. 21140). Seedless fruits were dried in a greenhouse with air circulation, powdered in a pulverizer mill and passed through a 60-mesh sieve at the Laboratory of Research on Natural Products, Pharmacy School/UFG. The powder obtained was stored at -20°C. To prepare the extract, the dried material was exhaustively percolated into an ethanol:water solution (55:45 v/v), and the material obtained was filtered and submitted to rotary evaporation under reduced pressure at 40°C, resulting in the ethanol-free *jaboticaba* hydroalcoholic extract (JHE). After that process, the JHE was maintained in a freezer (-20°C) protected from light. On the days of experiment, the JHE was solubilized in distilled water at the concentration of 120 mg/mL.

The phytochemical characterization and pattern of the JHE showed 17.89% of phenolic compounds, quantified by using the Hagerman and Butler method, adapted by Mole and Watermen.¹⁵ The JHE showed ellagic acid (phytochemical marker, determined by HPLC-PDA) at 0.222% concentration.¹⁴ According to Abe et al.,⁹ the total ellagic acid content in *M. cauliflora* fruits ranges from 0.021% to 0.311%. Thus, that phytochemical marker concentration in JHE is in accordance with the fruit content.

Animals and preparation of isolated arteries

Wistar male rats (200-230 g) from the central vivarium of the UFG were used in this study. All experimental protocols abided by the UFG Animal Research Ethics Committee (protocol: 015/2014). This study is in accordance with the European Union Guide to the Care and Use of Experimental Animals (2010/63/UE).

The rats were sacrificed by use of cervical dislocation under inhalation anesthesia. Thoracic aorta was isolated, cleared of connective and adipose tissues, and sliced into rings (\pm 4 mm), which were mounted between two metal hooks, one of which was connected to a power transducer to record isometric tension (DATAQ Instruments, Akron, OH, USA) and the other was fixed to a cube for the isolated organ. The rings were placed into chambers for isolated organs, containing modified Krebs solution [composition in mM: NaCl, 130.0; KCl, 4.7; KH_2PO_4 , 1.2; CaCl_2 , 1.6; MgSO_4 , 1.2; NaHCO_3 , 14.9; glucose, 5.5], at pH of 7.4, under gasification with carbogen mixture (95% O_2 + 5% CO_2) at 37°C, and maintained at baseline tension of 1 g (ideal resting tension, previously standardized at our laboratory). To prevent the influence of vascular-endothelium-derived factors, endothelial cells were mechanically removed by rubbing the vessel lumen with a thin metal rod, the effectiveness of the removal being evidenced by lack of relaxation due to acetylcholine (1 μM) in aortic rings pre-contracted with EC_{50} of phenylephrine (0.1 μM).

Experimental protocols

After 60 minutes of stabilization at baseline tension (1 g), the arteries were pre-contracted with phenylephrine (0.1 μM), and cumulative relaxation-concentration-effect curves were built for JHE (0 to 1.92 mg/mL) and for verapamil, used as inner control (10 nM to 100 μM).

To assess the cellular pathways responsible for the relaxant effect of JHE, aortic rings were pre-contracted with phenylephrine (0.1 μM) for 20 minutes after incubation with the following agents: 1) Ca^{2+} ATPase of the sarcoplasmic reticulum, cyclopiazonic acid (CPA, 10 μM); 2) non-selective K^+ channel blocker, tetraethyl-ammonium (TEA, 1 mM); 3) selective voltage-gated K^+ channel (K_v) blocker, 4-aminopyridine (4-AP, 1 mM); 4) selective ATP-sensitive K^+ channel (K_{ATP}) blocker, glibenclamide (3 μM); 5) Ca^{2+} -dependent K^+ channel (K_{Ca}) blocker, clotrimazole (5 μM).

To assess the influence of JHE on the contraction induced by adrenergic contractile agonist, concentration-effect curves were built for phenylephrine (selective α 1-adrenergic agonist, 0.1 nM to 10 μM) in the presence (20 minutes) or absence of JHE at inhibitory concentration 50% (IC_{50} , 0.5 mg/mL) or 100% (IC_{100} , 1.92 mg/mL). In addition, the inhibitory effect of the Ca^{2+} channel blocker verapamil (IC_{50} , 0.3 μM) was assessed as inner control.

In another series of experiments, JHE action on Ca^{2+} influx stimulated by two different agents was analyzed. The preparations were initially contracted with a KCl solution (75 mM) to cause maximum contraction of each preparation (100% contraction), being then rinsed with Ca^{2+} -free Krebs solution until total relaxation. To exhaust the intracellular storage of Ca^{2+} , the preparations were stimulated to contract with phenylephrine in Ca^{2+} -free Krebs solution until any contractile response disappeared (approximately 5 or 6 times, for 30-50 minutes). Then the preparations were rinsed several times with Ca^{2+} -free Krebs solution, and then incubated for 20 minutes with JHE at inhibitory concentration 50% (IC_{50} , 0.51 mg/mL) or 100% (IC_{100} , 1.92 mg/mL). In addition, the inhibitory effect of the Ca^{2+} channel blocker verapamil (IC_{50} , 0.3 μM) was assessed as inner control. After incubation, the contractile stimulus was applied (phenylephrine, 0.1 μM , or KCL, 75 mM), and concentration-effect curves were built for CaCl_2 (0 to 3.0 mM).

Statistical analysis

The results of isometric tension were expressed as mean \pm standard error of the mean (SEM) of at least five experiments ($n = 5-8$) obtained from different animals. The graphs were built and analyzed by use of the GraphPad Prism software (GraphPad Software Corporation, 5.0 version) with ANOVA and Bonferroni post-test. The 5% significance level ($p < 0.05$) was adopted for the differences.

Results

Relaxant effect of JHE on isolated arteries

The JHE caused relaxation in preparations of endothelium-denuded arteries on a concentration-dependent way,

relaxation initiating at the concentration of 0.38 mg/mL, and achieving the maximum effect (E_{max}) of $98.3\% \pm 0.4\%$ ($n = 6$) at the concentration of 1.92 mg/mL (IC_{100}) (Figure 1A). The JHE concentration that induced 50% relaxation (IC_{50}) was 0.51 mg/mL. Similarly, verapamil (used as positive control) induced concentration-dependent relaxation with E_{max} of $99.8\% \pm 1.8\%$ ($n = 5$) and IC_{50} of $0.3 \mu M$.

Effect of JHE on the phenylephrine-induced contraction

The E_{max} value for phenylephrine ($142.1\% \pm 7.1\%$, $n = 6$) was significantly ($p < 0.001$) reduced to $88.7\% \pm 6.2\%$ ($n = 5$), $66.1\% \pm 5.1\%$ ($n = 6$) and $79.9\% \pm 5.5\%$ ($n = 5$) after incubation with IC_{50} and IC_{100} of JHE or verapamil, respectively. The addition of IC_{50} and IC_{100} of JHE or verapamil significantly increased phenylephrine pD_2 values ($-\log EC_{50}$) from 6.24 ± 0.09 to 5.35 ± 0.04 , 5.14 ± 0.09 and 5.68 ± 0.07 , respectively (Figure 2).

Effect of JHE on Ca^{2+} -influx-induced contraction in preparations stimulated with phenylephrine or KCl

Regarding the Ca^{2+} -influx-induced contraction stimulated by phenylephrine, pre-incubation with JHE (IC_{50} or IC_{100}) significantly reduced ($p < 0.001$) the E_{max} values from $106.8\% \pm 7.5\%$ ($n = 5$) to $58.8\% \pm 4.9\%$ ($n = 6$) and $34.5\% \pm 3.2\%$ ($n = 6$), respectively. In addition, treatment

with verapamil significantly reduced ($p < 0.001$) the contraction to $7.1\% \pm 1.1\%$ ($n = 5$) (Figure 3A).

Regarding the Ca^{2+} -influx-induced contraction stimulated by KCl (75 mM), pre-incubation with JHE (IC_{50} or IC_{100}) significantly reduced ($p < 0.001$) the E_{max} values from $108.8\% \pm 4.3\%$ ($n = 5$) to $63.8\% \pm 6.1\%$ ($n = 6$) and $14.6\% \pm 1.9\%$ ($n = 6$), respectively. In addition, treatment with verapamil significantly reduced ($p < 0.001$) the contraction to $15.5\% \pm 1.1\%$ ($n = 6$) (Figure 3B).

Effect of reticular Ca^{2+} ATPase inhibitor, CPA, and K^+ -channel blockers on JHE-induced relaxation

Treatment with CPA did not change the JHE-induced relaxation ($93.8\% \pm 4.6\%$, $n = 6$) in isolated arteries (Figure 4). Thus, JHE did not change the inner Ca^{2+} uptake by the sarcoplasmic reticulum to induce vascular relaxation.

As shown in figure 5, except for clotrimazole ($94.1\% \pm 4.5\%$, $n = 5$), K^+ -channel blockers changed the JHE-stimulated relaxation. The JHE-induced relaxation (E_{max} : $98.3\% \pm 0.4\%$, $n = 6$) was significantly ($p < 0.05$) reduced by TEA (E_{max} : $87.6\% \pm 5.7\%$, $n = 5$), glibenclamide (E_{max} : $61.6\% \pm 5.8\%$, $n = 6$) and 4-AP (E_{max} : $81.6\% \pm 5.9\%$, $n = 5$). The results showed that JHE-induced relaxation depends on K^+ efflux through the membrane.




Figure 1 – Cumulative concentration-response curves of jaboticaba hydroalcoholic extract (JHE) (A) and verapamil (B) in isolated endothelium-denuded arteries. The points represent mean \pm SEM of the relaxant effect expressed as % relaxation.




Figure 2 – Effect of jaboticaba hydroalcoholic extract (JHE) and verapamil on the phenylephrine-induced contraction in isolated endothelium-denuded arteries. Cumulative concentration-response curves were built in control conditions and after incubation (20 min) with JHE (IC_{50} : 0.51 or IC_{100} : 1.92 mg/mL) or verapamil (IC_{50} : 0.3 μ M). The points represent mean \pm SEM of the contractile effect expressed as % contraction in relation to total KCl-induced contraction (75 mM). Significant difference: *** $p < 0.001$ vs. Control.




Figure 3 – Effect of jaboticaba hydroalcoholic extract (JHE) and verapamil on the contractile response in isolated endothelium-denuded arteries. Cumulative concentration-response curves for $CaCl_2$ were stimulated with phenylephrine (0.1 μ M) (A) or KCl 75 mM (B) in control conditions and after incubation (20 min) with JHE (IC_{50} : 0.51 or IC_{100} : 1.92 mg/mL) or verapamil (IC_{50} : 0.3 μ M). The points represent mean \pm SEM of the contractile effect expressed as % contraction in relation to total KCl-induced contraction (75 mM). Significant difference: *** $p < 0.001$ vs. Control.

Discussion

The major finding of this study is that JHE, in addition to having a hypotensive effect and inducing vascular relaxation through endothelial nitric oxide pathway, as shown by our team,¹⁴ acts directly on VSM and leads to endothelium-independent relaxation. Therefore, *jaboticaba* clearly has

cardiovascular effects through multiple endothelium-dependent and independent pathways. It is worth noting that the JHE concentration capable of inducing 100% vascular relaxation through the endothelial pathway is approximately 16 times lower (0.12 mg/mL)¹⁴ than the JHE concentration necessary to induce 100% relaxation acting directly on VSM (1.92 mg/mL).




Figure 4 – Maximum relaxant effect induced by jaboticaba hydroalcoholic extract (JHE) in isolated arteries pre-contracted with phenylephrine (0.1 μ M) in the absence or presence (20 min) of the reticular Ca^{2+} ATPase inhibitor, cyclopiazonic acid (CPA, 10 μ M). The bars represent mean \pm SEM of the maximum relaxant effect expressed as % relaxation.




Figure 5 – Effects of K^+ channel blockers on the relaxation induced by jaboticaba hydroalcoholic extract (JHE) in isolated arteries pre-contracted with phenylephrine (0.1 μ M) in the absence or presence (20 min) of the blockers tetraethyl-ammonium (TEA, 1 mM), glibenclamide (Glib, 3 μ M), clotrimazole (5 μ M) and 4-aminopyridine (4-AP, 1 mM). The points represent mean \pm SEM of the relaxant effect expressed as % relaxation. Significant difference: * $p < 0.05$; *** $p < 0.001$ vs. Control

Blood vessel contraction and relaxation in response to physiological demands are controlled by changes in intracellular Ca^{2+} concentration of VSM. The Ca^{2+} used for contraction includes intracellular or extracellular sources, or both. Sarcoplasmic reticulum is the major source of intracellular Ca^{2+} .¹⁶ Our experiments showed that JHE does not change Ca^{2+} uptake by the sarcoplasmic reticulum, because its selective inhibitor, CPA, did not change the relaxation profile.

Voltage-gated Ca^{2+} channels (VGCC), also known as L-type Ca^{2+} channels, and receptor-operated Ca^{2+} channels (ROCC) located on the plasma membrane of VSM cells play a fundamental role in controlling Ca^{2+} influx.^{17,18} Phenylephrine-induced contraction is mediated by Ca^{2+} influx increase via VGCC and ROCC.^{19,20} However, contraction induced by membrane depolarization, such as in high KCl

concentrations, activates preferentially VGCC.²¹ The results of the present study show that treating arteries with JHE inhibits the vascular contraction induced by the adrenergic stimulus with phenylephrine, suggesting that JHE blocks Ca^{2+} influx by interfering with VGCC and/or ROCC.

In an attempt to clarify the cell mechanism through which JHE induces vascular relaxation, experiments were performed in a Ca^{2+} -free solution. Two different stimuli, phenylephrine and KCl (75 mM), were used to induce Ca^{2+} influx. The JHE, as well as verapamil, used as a positive control, inhibited the Ca^{2+} -influx-induced contraction mediated by both stimuli. Because membrane depolarization with high concentrations of K^+ activates specifically VGCC, we suggest that JHE acts directly or indirectly by blocking Ca^{2+} influx through the plasma membrane, acting preferentially on VGCC.

Natural products have constantly shown the involvement of K⁺ channels in their vasodilating mechanism.²² Several types of K⁺ channels, such as ATP-sensitive K⁺ channels (K_{ATP}), Ca²⁺-dependent K⁺ channels (K_{ca}) and voltage-gated K⁺ channels (K_v), are present in VSM.^{23,24} Those channels can be blocked by glibenclamide, clotrimazole and 4-AP, respectively.^{24,25} Tetraethyl-ammonium is a non-selective blocker of those channels. When activated, those channels allow K⁺ efflux, hyperpolarizing the VSM plasma membrane. This reduces Ca²⁺ influx through the VGCC and induces vasodilation.^{26,27} The present study shows that JHE-induced relaxation in endothelium-denuded arteries is hindered after K⁺ channel blockade. Except for clotrimazole, the other blockers hindered vascular relaxation, allowing relating its activation to the JHE effect.

Our results point to a new biological effect of *jaboticaba*, a Brazilian native specimen that has important biological effects on the cardiovascular system, such as glucose-lowering,¹² lipid-lowering¹³ and hypotensive effects.¹⁴ Thus, the biological *jaboticaba*-induced effects demonstrated in this study will contribute to increase the knowledge about *jaboticaba*-derived compounds and their use as medicinal plant or functional food to prevent cardiovascular problems.

Conclusion

This study shows that JHE induces endothelium-independent vasodilation. The major cellular pathways used by JHE to cause vascular relaxation are inhibition of the Ca²⁺-influx through plasma membrane, in addition to K⁺ channel activation in VSM cells.

References

1. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the global burden of disease study 2010. *Lancet*. 2012;380(9859):2224–60. Erratum in: *Lancet*. 2013;381(9867):628, *Lancet*. 2013;381(9874):1276.
2. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005;365(9455):217–23.
3. Carafoli E. Intracellular calcium homeostasis. *Annu Rev Biochem*. 1987;56:395–433.
4. Somlyo AV, Bond M, Somlyo AP, Scarpa A. Inositol trisphosphate-induced calcium release and contraction in vascular smooth muscle. *Proc Natl Acad Sci USA*. 1985;82(15):5231–5.
5. Baker EH. Ion channels and the control of blood pressure. *Br J Clin Pharmacol*. 2000;49(3):185–98.
6. Greenway F, Liu Z, Yu Y, Gupta A. A clinical trial testing the safety and efficacy of a Standardized *Eucommia ulmoides* olive bark extract to treat hypertension. *Altern Med Rev*. 2011;16(4):338–47.
7. Chan P, Tomlinson B, Chen YJ, Liu JC, Hsieh MH, Cheng JT. A double-blind placebo-controlled study of the effectiveness and tolerability of oral stevioside in human hypertension. *Br J Clin Pharmacol*. 2000;50(3):215–20.
8. Fortes GA, Naves SS, Godoi FF, Duarte AR, Ferri PH, Santos SC. Assessment of a maturity index in jaboticaba fruit by the evaluation of phenolic compounds, essential oil components, sugar content and total acidity. *Am J Food Technol*. 2011;6(11):974–84.
9. Abe LT, Lajolo FM, Genovese MI. Potential dietary sources of ellagic acid and other antioxidants among fruits consumed in Brazil: jaboticaba (*Myrciaria jaboticaba* (Vell.) Berg). *J Sci Food Agric*. 2012;92(8):1679–87.
10. Giraldo M, Hanazaki N. Use and traditional knowledge of medicinal plants at Sertão do Ribeirão, Florianópolis, Santa Catarina State, Brazil. *Acta Bot Bras*. 2010;24(2):395–406.
11. Reynertson KA, Yang H, Jiang B, Basile MJ, Kennelly EJ. Quantitative analysis of antiradical phenolic constituents from fourteen edible Myrtaceae fruits. *Food Chem*. 2008;109(4):883–908.
12. Dragano NR, Marques AY, Cintra DE, Solon C, Morari J, Leite-Legatti AV, et al. Freeze-dried jaboticaba peel powder improves insulin sensitivity in high-fat-fed mice. *Br J Nutr*. 2013;110(3):447–55.
13. Araújo CR, Esteves EA, Dessimoni-Pinto NA, Batista AG. *Myrciaria cauliflora* peel flour had a hypolipidemic effect in rats fed a moderately high-fat diet. *J Med Food*. 2014;17(2):262–7.
14. Lobo de Andrade DM, Reis CF, Castro PF, Borges LL, Amaral NO, Torres IM, et al. Vasorelaxant and Hypotensive Effects of Jaboticaba Fruit (*Myrciaria cauliflora*) Extract in Rats. *Evid Based Complement Alternat Med*. 2015;2015:696135.

Acknowledgements

This research was funded by the Goiás State Research Support Foundation (FAPEG), process 201310267000013. The Brazilian National Council for Scientific and Technological Development (CNPq) provided funding to support the Master's program of Daniela L. M. de Andrade.

Author contributions

Conception and design of the research: Andrade DML, Borges LL, Torres IMS, Conceição EC, Rocha ML; Acquisition of data: Andrade DML, Borges LL, Conceição EC, Rocha ML; Analysis and interpretation of the data: Andrade DML, Torres IMS, Conceição EC, Rocha ML; Statistical analysis: Andrade DML, Rocha ML; Obtaining financing: Rocha ML; Writing of the manuscript: Andrade DML, Torres IMS, Rocha ML; Critical revision of the manuscript for intellectual content: Andrade DML, Borges LL, Conceição EC, Rocha ML.

Potential Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Sources of Funding

This study was partially funded by FAPEG/GO.

Study Association

This article is part of the thesis of master submitted by Daniela Medeiros Lobo Andrade, from Faculdade de Farmácia - Universidade Federal de Goiás.

15. Mole S, Waterman PG. A critical analysis of techniques for measuring tannins in ecological studies-II. Techniques for biochemically defining tannins. *Oecologia*. 1987;72(1):148-56.
16. Thomas AP, Bird GS, Hajoczky G, Robb-Gaspers LD, Putney JW Jr. Spatial and temporal aspects of cellular calcium signaling. *FASEB J*. 1996;10(13):1505-17.
17. Berridge MJ. Cell signalling: a tale of two messengers. *Nature*. 1993;365(6445):388-9.
18. Hess P. Calcium channels in vertebrate cells. *Annu Rev Neurosci*. 1990;13:337-56.
19. Wesselman JP, VanBavel E, Pfaffendorf M, Spaan JA. Voltage-operated calcium channels are essential for the myogenic responsiveness of cannulated rat mesenteric small arteries. *J Vasc Res*. 1996;33:32-41.
20. Lee CH, Poburko D, Sahota P, Sandhu J, Ruehlmann DO, van Breemen C. The mechanism of phenylephrine-mediated $[Ca^{2+}]_i$ oscillations underlying tonic contraction in the rabbit inferior vena cava. *J Physiol*. 2001;534(Pt 3):641-50.
21. Kochegarov AA. Pharmacological modulators of voltage-gated calcium channels and their therapeutic application. *Cell Calcium*. 2003;33(3):145-62.
22. McNeill JR, Jurgens TM. A systematic review of mechanisms by which natural products of plant origin evoke vasodilatation. *Can J Physiol Pharmacol*. 2006;84(8-9):803-21.
23. Ko EA, Han J, Jung ID, Park WS. Physiological roles of K^+ channels in vascular smooth muscle cells. *J Smooth Muscle Res*. 2008;44(2):65-81.
24. Ferrer M, Marin J, Encabo A, Alonso MJ, Balfagon G. Role of K^+ channels and sodium pump in the vasodilation induced by acetylcholine, nitric oxide, and cyclic GMP in the rabbit aorta. *Gen Pharmacol*. 1999;33(1):35-41.
25. Wulff H, Miller MJ, Hansel W, Grissmer S, Cahalan MD, Chandy KG. Design of a potent and selective inhibitor of the intermediate-conductance Ca^{2+} -activated K^+ channel, IK_{Ca1} : a potential immunosuppressant. *Proc Natl Acad Sci USA*. 2000;97(14):8151-6.
26. Nelson MT, Quayle JM. Physiological roles and properties of potassium channels in arterial smooth muscle. *Am J Physiol*. 1995;268(4 Pt 1):C799-822.
27. Ledoux J, Werner ME, Brayden JE, Nelson MT. Calcium activated potassium channels and the regulation of vascular tone. *Physiology (Bethesda)*. 2006;21:69-78.