

Effects of Acute Exercise on Postprandial Lipemia in Sedentary Men

Mônica Teixeira, Nelson Kasinski, Maria Cristina de Oliveira Izar, Luiz Alberto Barbosa, José Paulo Novazzi, Leonor Almeida Pinto, Sérgio Tufik, Turíbio Ferreira Leite, Francisco Antonio Helfenstein Fonseca

Universidade Federal de São Paulo – Escola Paulista de Medicina - São Paulo, SP - Brazil

OBJECTIVE

To examine the effects of a single session of isometric physical exercise on postprandial triglyceridemia in sedentary male individuals with fasting triglycerides values < 150 mg/dl (NTG) or ≥ 150 mg/dl (TG_{ALT}).

METHODS

Twenty-seven individuals (10 NTG and 17 TG_{ALT}), aged between 30-55 years were assessed in the study. Triglycerides were determined in the beginning, and at two, four and six hours after the oral ingestion of a solution with $50g/m^2$ of fat in two occasions: at rest and after treadmill isometric exercise.

RESULTS

Acute exercise did not affect the levels of postprandial triglycerides or the area under the curve (AUC) of triglycerides. However, the abnormal pattern of postprandial lipemia curve was associated with higher basal triglyceridemia with exercise (basal TG: 147 ± 90 vs. 238 ± 89 mg/dl, $p = 0.02$) and without exercise (basal TG: 168 ± 93 vs. 265 ± 140 mg/dl, $p = 0.04$). Analysis of the receiver operating characteristics (ROC) curves showed cut-off values for basal triglycerides with exercise of 166.5 mg/dl (sensitivity: 0.78; specificity: 0.72) and AUC of 0.772 [CI 95%: 0.588-0.955], and without exercise of 172 mg/dl (sensitivity: 0.78; specificity: 0.61) and AUC: 0.722 [CI 95%: 0.530-0.914].

CONCLUSION

Acute exercise did not affect postprandial triglyceridemia in sedentary male individuals, and basal triglyceride levels are predictors of an abnormal response of postprandial triglycerides.

KEY WORDS

Acute exercise, triglycerides, postprandial lipemia.

The concept that the atherosclerotic process is a postprandial phenomenon was evidenced more than twenty years ago by Zilversmit¹, who demonstrated, in animals and in human subjects, that remnants of chylomicrons and low-density lipoproteins (LDL) are captured by endothelial cells.

Clinical studies have shown that the magnitude of postprandial lipemia or even increased postprandial triglyceride (TG) levels can predict symptomatic or asymptomatic atherosclerosis, regardless of risk factors assessed in the fasting state²⁻⁵.

Postprandial lipemia reflects an integrated measure of the individual's capacity to remove triglycerides. Diets or treatments which reduce the fasting levels of triglyceride-rich lipoproteins may improve the postprandial levels of triglyceride-rich lipoproteins. Therefore, aerobic exercise, the reduction of body weight and certain medicines reduce postprandial triglyceridemia and can also reduce remnant lipoprotein levels⁶.

Postprandial lipemia is characterized by transient hypertriglyceridemia, but fasting triglyceride concentrations are related to the size of the LDL particle⁷. This response may be conditioned to the higher mobilization of fatty acids, increase of the synthesis and delay in the removal of VLDL, thus allowing greater interaction between lipoproteins which contribute to the formation of small and dense LDL particles which are extremely atherogenic⁷⁻¹⁰.

The magnitude of postprandial response is determined by several factors. Postprandial response is greater in men^{11,12}, tends to increase with age¹³, is different between normo- and hypertriglyceridemic individuals^{1,14} according to fasting serum lipid levels.

Regular exercise is associated with the reduction of the prevalence of coronary artery disease, diabetes mellitus and obesity¹⁵⁻¹⁸. Most of our life is spent in the postprandial state, and therefore interventions which may mitigate the atherogenic effects of postprandial hyperlipemia are specially important¹⁹.

Although exercise may cause favorable changes in postprandial lipid profile, the acute effects of exercise are less known, especially when individuals with normal or altered fasting triglycerides levels are compared²⁰⁻²².

In view of the considerable increase in the prevalence of sedentary lifestyle, obesity and insulin resistance in modern society, this study was designed to examine the effects of a single session of physical exercise on postprandial triglyceridemia, in sedentary individuals with or without fasting hypertriglyceridemia.

METHODS

The study was approved by the Research Ethics Committee of the Federal University of São Paulo

– Paulista School of Medicina, and all participants gave their written informed consent.

Twenty-seven sedentary individuals participated in the study, with ages between 30 and 55, of whom ten were normotriglyceridemic NTG (basal TG < 150 mg/dl) and seventeen had altered basal triglyceride values TG_{ALT} (TG ≥ 150)^{23,24}. Individuals with TG > 400 mg/dl were excluded from the study.

We also excluded from the study those individuals with body mass index ≥ 30 kg/m², with symptomatic coronary disease, assessed by means of clinical and electrocardiographic criteria, recorded at rest or during the treadmill exercise test. Additionally we excluded individuals with severe liver, severe kidney disease, diabetes, patients with infections or cancer.

Design of the study - The protocol comprised medical assessments, laboratory tests, ergospirometry, nutritional assessment and study of postprandial lipemia (fig. 1). Exercise prescription followed the ergospirometric test.

Nutritional assessment comprised two stages. We obtained the anthropometric data of the selected individuals in basal conditions, in addition to a 3-day feeding record²⁵. Based on the data informed, we calculated the usual diet. The ingestion of each nutrient in the usual diet was calculated with the Nutrition Support Software Program of the Center of Computers applied to Health – EPM/ UNIFESP – version 2.5²⁶.

The study of postprandial lipemia was carried out in two periods, with patients submitted to thirty minutes of exercise on the treadmill, right after the ingestion of the diet. The study was repeated two weeks later, without the treadmill exercise.


We collected blood samples after a twelve to fourteen hour fast and at two, four and six hours following the ingestion of the standardized diet for analysis of the triglycerides (fig. 1).

Laboratory analysis - Total cholesterol and triglycerides were measured using an automated enzymatic method (Opera, Bayer, Germany), the HDL-C fraction being quantified in the supernatant, after the precipitation of lipoproteins containing apo B; LDL-C was estimated using Friedewald's formula²⁷. Biochemical analyses were carried out in an Advia 120 device (Bayer, Germany).

Ergospirometric assessment and physical exercise - The ergospirometric test was carried out on a treadmill with a continuous protocol of increasing loads²⁸.

The exercise proposed was walking on the treadmill, with no slope, for thirty minutes at constant speed, according to the result of the ergospirometric test (anaerobic threshold). The exercise phase followed a 3-minute warm up phase, which was repeated at the end of each period.

Postprandial lipemia - The individuals were submitted to two tests for postprandial lipemia. As preparation prior to postprandial lipemia, they were asked to suspend



physical activity for 48 hours, to refrain from drinking alcohol and caffeine for a period of 24 hours, and to follow the diet prescribed for dinner in the evening before the test²⁹⁻³³.

Blood samples were collected using a plastic catheter (Vialon® n. 20, Becton Dickinson), and 5 ml of blood were discarded at each new sample collection to guarantee the precision of the test results. Four blood samples were collected (fasting, two, four and six hours). After the basal collection, the participants ingested a solution with a high concentration of fats, in fifteen minutes.

The solution was made up of: one liter of whole milk, one can of milk cream, 50 grams of dehydrated hen egg, 100 grams of powder chocolate, one tablespoon of saltless butter, six Brazil nuts, two dwarf bananas, 150 ml of coconut milk. The total nutritional value of the recipe was 2,655 calories, with 81 g of proteins (11.6%), 151 g of carbohydrates (21.4%) and 209 g of lipids (67.0%).

Each participant received the equivalent to 50 grams of fat per square meter of body surface. The yield of this recipe relative to 50 grams of fat was 365 ml which, multiplied by the body surface, resulted in the ingested quantity of the solution.

Statistical analysis - Numerical variables of the basal period are presented as means and standard deviation relative to the means and were compared across the NTG and TG_{ALT} groups using Student's *t* test, for unrelated samples. For the analysis of triglycerides in fasting, and then at two, four and six hours, we used the repeated measures analysis of variances. The area under the curve was calculated using the trapezoidal rule. We used Student's *t* test for related samples to compare the areas under the curves with and without physical activity. We built ROC curves to assess the efficiency of triglyceride basal values in the prediction of alterations in the response of postprandial triglyceridemia. Values of $p < 0.05$ were considered significant.

RESULTS

The basal characteristics of the participants are presented in Table 1. There was no difference in demographic variables and anthropometric parameters between NTG and TG_{ALT} participants, with the exception of waist circumference, which was larger in TG_{ALT} individuals. In addition to higher TG basal values, TG_{ALT} individuals presented lower HDL-C values when compared with NTG individuals, with no difference as regards glycemia values.

There was no difference across the groups as regards the total calorie consumption of the diet, the content of proteins, carbohydrates, and lipids; however, NTG individuals presented greater consumption of polyunsaturated fatty acids than TG_{ALT} individuals (Table 2).

Postprandial lipemia and the effect of acute exercise - Postprandial triglyceridemia peaked within four hours, with no difference as regards the prescription of acute exercise or not ($p = 0.098$), with a ratio between the values of triglycerides at peak of 0.95 (CI 95%: 0.74;1.15) (fig. 2). Mean values obtained for the areas under the triglyceride curve (six hours) did not differ (AUC: 632 ± 422 vs. 751 ± 427 mg/dl, with and without physical activity, respectively; $p = 0.133$, Student's *t* test for related samples), with the ratio of the areas under the curve being 0.97 (CI 95%:0.75;1.19) (fig. 3).

Postprandial triglyceride curves are presented in Table 3 and in figure 4. There was no effect of acute exercise on the postprandial triglyceride curve ($p = 0.40$; ANOVA with repeated measures). However, there was an association of the basal triglyceride values with the behavior of the triglyceride curve of individuals in time ($p = 0.026$).

Analyzing the behavior of the postprandial triglyceride curve of the participants of the study we observed that eighteen individuals presented a behavior which was typical of a normal curve, i.e., at hour six after the

Table 1 – Demographic and anthropometric characteristics and basal biochemical variables of NTG and TG_{ALT} participants

Variables	NTG N = 10	SD	TG _{ALT} N = 17	SD	P
Age (median, years)	41	-	42	-	0.990
BMI (Kg/m ²)	25	2	25	3	0.670
Waist circumference (cm)*	82	7	91	7	0.006
Total cholesterol (mg/dl)	204	29	227	37	0.110
HDL-C (mg/dl)*	52	12	38	11	0.007
LDL-C (mg/dl)	134	28	145	41	0.430
Triglycerides (mg/dl)**	110	18	259	68	< 0.0001
Glycemia (mg/dl)	93	8	96	11	0.430

SD = standard deviation relative to mean. *TG_{ALT} < NTG; p < 0.05; **TG_{ALT} > NTG; p < 0.05; Student's t test for independent samples.

Table 2 – Distribution of NTG and TG_{ALT} participants according to food consumption

Dietary variable	NTG	SD	TG _{ALT}	SD	p
Total calories (Kcal/d)	2195	590	1980	735	0.44
Proteins (%)	20	2	22	5	0.10
Carbohydrates (%)	44	10	48	8	0.26
Lipids (%)	35	8	30	6	0.06
Saturated fatty acids (%)	12	4	10	3	0.28
Polyunsaturated fatty acids (%)*	5	2	3	1	0.01
Monounsaturated fatty acids (%)	9	4	6	3	0.06
Cholesterol (mg)	331	109	285	103	0.29

SD = standard deviation relative to mean. *NTG > TG_{ALT}; p < 0.05; Student's t test for independent samples.




Fig. 2 – Boxplots of peak triglyceride values (4 hours) with and without physical activity (PA). PA = physical activity. With PA = Without PA; p = 0.098; Student's t test for related samples. Ratio between the values obtained in the peak = 0.95 (CI 95%: 0.74;1.15).




Fig. 3 – Boxplots of areas under curve of triglycerides values with and without physical activity. PA = physical activity. With PA = Without PA; p = 0.133; Student's t test for related samples. Ratio between the values obtained for the areas under the curves = 0.97 (CI 95%: 0.75; 1.19).

ingestion of the lipid solution triglyceridemia returned to near basal values, whereas nine subjects presented abnormal curves. Individual profiles of postprandial lipemia response with and without physical activity are presented in figures 5 and 6, respectively. Basal triglyceride values were then compared according to the behavior of the normal or abnormal postprandial triglyceride curve, with and without physical activity, with higher basal triglyceride values for individuals who had an

abnormal postprandial lipemia curve with physical activity (basal TG: 147 ± 90 vs 238 ± 89 mg/dl, p = 0.02, Student's t test) and without physical activity (basal TG: 168 ± 93 vs 265 ± 140 mg/dl, p = 0.04, Student's t test) (fig. 7 and 8). There was agreement in the response of postprandial triglyceridemia in seventeen individuals with and without acute exercise (thirteen normal with and without physical activity and four abnormal individuals with and without physical activity).

Table 3 – Distribution of triglyceride values throughout the postprandial lipemia curve with and without physical activity

	Beginning	SD	2h	SD	4h	SD	6h	SD
Without PA	201	23	289	27	406	41	372	55
With PA	177	19	258	24	340	36	325	48

PA = physical activity; SD = standard deviation relative to mean. With PA = without PA; p = 0.400; ANOVA with repeated measures. Beginning < 2h, 4h and 6h; p = 0.026, ANOVA with repeated measures.




Fig. 4 – Mean values of the responses of triglycerides in postprandial lipemia with and without physical activity. PA = physical activity. With PA = Without PA; p = 0.400; ANOVA with repeated measures.




Fig. 5 – Individual profiles in the response of postprandial triglyceridemia with physical activity.




Fig. 6 – Individual profiles in the response of postprandial triglyceridemia without physical activity.




Fig. 7 – Boxplots of basal triglyceride values with physical activity according to the behavior of individuals in the postprandial triglyceride curve. PA = physical activity. Abnormal > Normal; $p = 0.020$; Student's t test for unrelated samples.




Fig. 8 – Boxplots of basal triglyceride values without physical activity according to the behavior of individuals in the postprandial triglyceride curve. PA = physical activity. Abnormal > Normal; $p = 0.040$; Student's t test for unrelated samples.

ROC curves showed that triglyceride basal values were predictors of an altered response of postprandial triglyceridemia with and without physical activity (fig. 9 and 10). The efficiency of basal triglycerides in predicting an altered response of postprandial triglycerides was 0.793 (CI 95%: 0.622-0.964), with physical activity and 0.722 (CI 95%: 0.530-0.914) without physical activity.

The cut-off point of triglyceride basal values for predicting an abnormal response of postprandial triglyceridemia was 166.5 mg/dl (sensitivity = 0.77 and specificity = 0.72) with physical activity and 172 mg/dl (sensitivity = 0.78 and specificity = 0.61) without physical activity.

DISCUSSION

The major contribution of this study was to show that acute physical exercise was unable to modify postprandial

triglyceridemia in sedentary men, and especially of valuing fasting triglyceride serum levels as predictors of postprandial lipemia. Additionally, it identified which triglyceride basal values are associated with abnormal postprandial triglyceridemia.

The study has also shown that sedentary individuals with altered fasting triglyceride levels (TG_{ALT}) presented larger waist circumference and lower HDL-C, although their mean BMI was similar to the BMI of normotriglyceridemic individuals.

Although the postprandial assessment of triglycerides did not evidence the effect of physical exercise on postprandial lipemia, we observed that a group of individuals showed an abnormal pattern of response of postprandial triglyceride, with or without physical activity, which characterized an abnormal behavior of the curve, and that this pattern could be predicted based on basal triglyceride values. Basal triglyceride values above 166.5 and 172.0 mg/dl, with and without physical activity,




Fig. 9 – ROC curve of basal triglyceride values with physical activity. Area under the curve: 0.772 (CI 95%: 0.588; 0.955). Basal triglyceride cut-off value: 166.5 mg/dl. Sensitivity: 0.78; Specificity: 0.72.




Fig. 10 – ROC curve of basal triglyceride values without physical activity. Area under the curve: 0.722 (CI 95%: 0.530; 0.914). Basal triglyceride cut-off value: 172 mg/dl. Sensitivity: 0.78; Specificity: 0.61.

respectively, efficiently predicted an altered response of postprandial triglycerides in sedentary individuals.

Triglyceride-rich lipoproteins from the intestine and liver contribute to the triglyceridemia found after the ingestion of a fat-rich solution^{2,10}. The quantification of lipemia was based on triglyceride serum concentration, which reflects the balance between the rate of appearance in and disappearance from the circulation, of lipoproteins rich in this lipid²⁹, with the peak of the lipemia curve being obtained four hours after the ingestion of the fatty solution¹⁴, and its normalization occurring as of the sixth hour³¹.

It is well established that during prolonged and marked postprandial lipemia there is exchange of neutral lipids between triglyceride-rich particles, LDL and HDL, thus leading to the formation of small and dense LDL particles and to a lower concentration of HDL-C³⁴. More recently, there were reports stating that endothelial function is altered in the postprandial state and is related to the increase of postprandial triglycerides³⁵. Other studies have also shown that, in healthy individuals, serum concentrations of inflammatory cytokines and adherence molecules were elevated after the ingestion of diets rich in fat, but not in carbohydrates³⁶, which caused a transitory proinflammatory state. These mechanisms are involved in the atherogenic process and are associated with greater cardiovascular risk³⁷.

Athletes and individuals who exercise regularly present low levels of postprandial lipemia³⁸. The effects of acute exercise in normotriglyceridemic individuals have been recently assessed, and no difference was observed in the response of postprandial triglycerides when the fat-rich diet was administered either before

or after the exercise³⁹. We assessed the effect of one week of detraining on the postprandial lipemia of trained individuals, and they presented increase in basal and postprandial triglycerides and of triglyceride-rich lipoproteins, with no change in the endothelial function in the short period without exercise⁴⁰.

In athletes, different degrees of acute exercise did not demonstrate changes in the area under the curves of postprandial triglycerides⁴¹. We did not find in the literature description of the effects of acute exercise on the postprandial response of triglycerides in hypertriglyceridemic individuals. In our study, with untrained normo- and hypertriglyceridemic individuals we did not observe any effect of a single session of exercise on postprandial triglyceridemia. However, basal triglyceride values were predictors of the postprandial response with or without acute exercise, thus reinforcing the orientation provided in current guidelines which emphasize the value of fasting lipid levels.

To conclude, our study evidenced that in untrained men, with normal or altered basal triglycerides values, acute exercise does not alter postprandial triglyceridemia. Basal triglyceride values above desirable ones are predictors of an abnormal response of postprandial triglycerides.

Acknowledge

We thank Prof. Dr. Fábio Tadeu Montesano for his assistance with the statistical analyses of this study.

Potencial Conflict of Interest

No potential conflict of interest relevant to this article was reported.

REFERENCES

1. Zilversmit DB. Atherogenesis: a postprandial phenomenon. *Circulation*. 1979; 60: 473-85.
2. Sharrett AR, Chambless LE, Heiss G, Paton CC, Patsch W, ARIC Investigators. Association of postprandial triglyceride and retinyl palmitate responses with asymptomatic carotid artery atherosclerosis in middle-aged men and women. *Arterioscler Thromb Vasc Biol*. 1995; 15: 2122-29.
3. Boquist S, Ruotolo G, Tang R, Björkegren J, Bond MG, De Faire U, et al. Alimentary lipemia, postprandial triglyceride-rich lipoproteins, and common carotid intima-media thickness in healthy, middle-aged men. *Circulation*. 1999; 100: 723-28.
4. Simons LA. Chylomicrons and chylomicron remnants in coronary artery disease: a case-control study. *Atherosclerosis*. 1987; 65: 181-89.
5. Groot PHE, van Stiphout WAHJ, Krauss XH. Postprandial lipoprotein metabolism in normolipidemic men with and without coronary artery disease. *Arterioscler Thromb*. 1991; 11: 653-62.
6. Cohn JS. Postprandial lipemia: emerging evidence for atherogenicity of remnant lipoproteins. *Can J Cardiol*. 1998; 14 Suppl B: 18B-27B.
7. Lemieux I, Couillard C, Pascot A, Bergeron N, Prud'homme D, Bergeron J, et al. The small, dense LDL phenotype as a correlate of postprandial lipemia in men. *Atherosclerosis*. 2000; 153: 423-32.
8. Austin MA, Breslow JL, Hennekens CH, Buring JE, Willett WC, Krauss RM. Low density lipoprotein subclass patterns and risk of myocardial infarction. *JAMA*. 1988; 260: 1917-21.
9. Ebenbichler CF, Kirchmair R, Egger C, Patsch JR. Postprandial state and atherosclerosis. *Curr Opin Lipidol*. 1995; 6: 286-90.
10. Karpe F, Tornvall P, Olivecrona T, Steiner G, Carlson LA, Hamsten A. Composition of human low density lipoprotein: effects of postprandial triglyceride-rich lipoproteins, lipoprotein lipase, hepatic lipase and cholesteryl ester transfer protein. *Atherosclerosis*. 1993; 98: 33-49.
11. Kashyap ML, Barnhart RL, Srivastava S, Perisutti G, Allen C, Hogg E, et al. Alimentary lipemia: plasma high-density lipoproteins and apolipoproteins C-II and C-III in healthy subjects. *Am J Clin Nutr*. 1983; 37: 233-43.
12. Roche HM, Gibney MJ. Effect of long-chain n-3 polyunsaturated fatty acids on fasting and postprandial triacylglycerol metabolism. *Am J Clin Nutr*. 2000; 7 (suppl): S232-7.
13. Krasinski SD, Cohn JS, Schaefer EJ, Russel RM. Postprandial plasma retinyl ester response is greater in older subjects compared with younger subjects. Evidence for delayed plasma clearance of intestinal lipoproteins. *J Clin Invest*. 1990; 85: 883-92.
14. Aldred HE, Perry IC, Hardman AE. The effect of a single bout of brisk walking on postprandial lipemia in normolipidemic young adults. *Metabolism*. 1994; 43: 836-41.
15. US Department of Health and Human Services. Physical activity and health: a report of the US Surgeon General. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion; 1996.

16. Berlin JA, Colditz GA. A meta-analysis of physical activity in the prevention of coronary heart disease. *Am J Epidemiol.* 1990; 132: 612-28.
17. Harris GD, White RD. Lifestyle modifications for the prevention and treatment of cardiovascular disease: an evidence-based approach. *Mo Med.* 2004; 101: 222-6.
18. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med.* 2002; 346: 393-403.
19. Blackburn P, Cote M, Lamarche B, Couillard C, Pascot A, Tremblay A, et al. Impact of postprandial variation in triglyceridemia on low-density lipoprotein particle size. *Metabolism.* 2003; 52: 1379-86.
20. Crouse SF, O'Brien BC, Grandjean PW, Lowe RC, Rohack JJ, Green JS. Effects of training and a single session of exercise on lipids and apolipoproteins in hypercholesterolemic men. *J Appl Physiol.* 1997; 83: 2019-28.
21. Ferguson MA, Alderson NL, Trost SG, Essig DA, Burke JR, Durstine L. Effects of four different single exercise sessions on lipids, lipoproteins, and lipoprotein lipase. *J Appl Physiol.* 1998; 85: 1169-74.
22. Hambrecht R, Niebauer J, Marburger C, Grunze M, Kälberer B, Haver K, et al. Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *J Am Coll Cardiol.* 1993; 22: 468-70.
23. III Diretrizes Brasileiras sobre Dislipidemias e Diretriz de Prevenção da Aterosclerose do Departamento de Aterosclerose. *Arq Bras Cardiol.* 2001; 77 (supl): 1-48.
24. Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults. Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA.* 2001; 285: 2486-96.
25. Basiotis PP. Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *J Nutr.* 1987; 117: 1138-46.
26. Anção NS, Cuppari L, Tudisco ES, Draibe SA, Sigulen D. Sistema de Apoio a decisão em Nutrição- versão 2,5 1995. Centro de Informática em Saúde- CIS- Universidade Federal de São Paulo-UNIFESP/EPM.
27. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge. *Clin Chem.* 1972; 18: 499-502.
28. Buchfehrer MJ, Hansen JE, Robson TE. Optimizing the exercise protocol for cardiopulmonary assessment. *J Appl Physiol.* 1983; 55: 1558-64.
29. Ziogas GG, Thomas TR, Harris WS. Exercise training, postprandial hypertriglyceridemia, and LDL subfraction distribution. *Med Sci Sports Exerc.* 1997; 29: 986-91.
30. Karpe F, Tornvall P, Olivecrona T, Steiner G, Carlson LA, Hamsten A. Composition of human low density lipoprotein: effects of postprandial triglyceride-rich lipoproteins, lipoprotein lipase, hepatic lipase and cholesteryl ester transfer protein. *Atherosclerosis.* 1993; 98: 33-49.
31. Tsetsonis NV, Hardman AE. Effects of low and moderate intensity treadmill walking on postprandial lipaemia in healthy young adults. *Eur J Appl Physiol.* 1996; 73: 419-26.
32. Hardman AE. The influence of exercise on postprandial triacylglycerol metabolism. *Atherosclerosis.* 1998; 141 (suppl 1): S93-100.
33. Hughes AT. Postprandial lipoprotein responses in hypertriglyceridemic subjects with and without cardiovascular disease. *Metabolism.* 1995; 44 (8): 1082-98.
34. Miesenböck G, Patsch JR. Postprandial hyperlipidemia: the search for the atherogenic lipoprotein. *Curr Opin Lipidol.* 1992; 3: 196-201.
35. Marchesi S, Lupatelli G, Schillaci G, et al. Impaired flow-mediated vasoactivity during postprandial phase in young healthy men. *Atherosclerosis.* 2000; 153: 397-402.
36. Nappo F, Espósito K, Cioffi M, et al. Postprandial endothelial activation in healthy subjects and in type 2 diabetic patients: role of fat and carbohydrate meals. *J Am Coll Cardiol.* 2002; 39: 145-50.
37. Plutzky J. Inflammatory pathways in atherosclerosis and acute coronary syndromes. *Am J Cardiol.* 2001; 88: 10K-15K.
38. Hardman AE, Lawrence JEM, Herd SL. Postprandial lipaemia in endurance-trained people during a short interruption to training. *J Appl Physiol.* 1998; 84: 1895-901.
39. Katsanos CS, Moffatt RJ. Acute effects of premeal versus postmeal exercise on postprandial hypertriglyceridemia. *Clin J Sport Med.* 2004; 14: 33-9.
40. Gill JMR, Caslake MJ, McAllister C, Tsofliou F, Ferrell WR, Packard CJ, et al. Effects of short-term detraining on postprandial metabolism, endothelial function, and inflammation in endurance-trained men: dissociation between changes in triglyceride metabolism and endothelial function. *J Clin Endocrinol Metab.* 2003; 88: 438-35.
41. Shannon KA, Shannon RM, Clore JN, Gennings C, Warren BJ, Potteiger JA. Resistance exercise and postprandial lipemia: The dose effect of differing volumes of acute resistance exercise bouts. *Metabolism.* 2005; 54: 756-63.